


ANNEX D - BIBLIOGRAPHY

Understanding the combined biodiversity benefits of the component features of hedges

Defra: Contract number BD5214

Robert J. Wolton, Roger K.A. Morris, Katie A. Pollard & John W. Dover

28 March 2013



Contract title: **Understanding the combined biodiversity benefits of the component features of hedges**

Client: Defra: Contract number BD5214

Prepared by Robert J. Wolton¹, Roger K.A. Morris², Katie A Pollard³ & John W. Dover⁴

¹Locks Park Farm

²Bright Angel Coastal Consultants Ltd

³KP Ecology Ltd

⁴Staffordshire University

Bright Angel Coastal Consultants Ltd.

7 Vine Street

Stamford

Lincolnshire

PE9 1QE

01780-753419

brightangel.coastal@googlemail.com

www.bacoastal.co.uk

Recommended citation:

Wolton, R.J., Morris, R.K.A., Pollard, K.A. & Dover J.W., 2013. Understanding the combined biodiversity benefits of the component features of hedges. Report of Defra project BD5214. Annex D: Bibliography.

BIBLIOGRAPHY

- AKSOY, S., AKÇAY, H. G., & WASSENAAR, T. 2010. Automatic mapping of linear woody vegetation features in agricultural landscapes using very high resolution imagery. *IEEE Transactions on Geoscience and Remote Sensing*, 48: 511-522.
- AL HASSAN, D., GEORGELIN, E., DELATTRE, T., BUREL, F., PLANTEGENEST, M., KINDLMANN, P., & BUTET, A. 2013. Does the presence of grassy strips and landscape grain affect the spatial distribution of aphids and their carabid predators? *Agricultural and Forest Entomology*, 15: 24–33.
- ALEXANDER, K.N.A. 2002. The invertebrates of living and decaying timber in Britain and Ireland. *English Nature Research Reports No 47*. English Nature, Peterborough. 142pp.
- ALEXANDER, K.N.A., 2004. Landscapes with ancient trees: invertebrate mobility and population viability. In: SMITHERS, R. (ed). *Landscape Ecology of Trees and Forests: Proceedings of the twelfth annual IALE (UK) conference, held at the Royal Agricultural College, Cirencester, 21st-24th June 2004* pp. 107-114.
- AQUILINA, R., WILLIAMS P. & NICOLET P. 2007. Effect of wetting-up ditches on emergent insect numbers. *Aspects of Applied Biology*, 81: 261-262.
- ARNOLD, G. W. 1983. The influence of ditch and hedgerow structure, length of hedgerows, and area of woodland and garden on bird numbers on farmland. *Journal of Applied Ecology*, 20: 731-750.
- AVIRON, S., BUREL, F., BAUDRY, J., & SCHERMANN, N. 2005. Carabid assemblages in agricultural landscapes: Impacts of habitat features, landscape context at different spatial scales and farming intensity. *Agriculture, Ecosystems and Environment*, 108: 205-217.
- BALENT, G., & COURTIADE, B. 1992. Modelling bird communities/landscape patterns relationships in a rural area of south-western France. *Landscape Ecology*, 6: 195-211.
- BARR, C. J. & GILLESPIE, M.K. 2000. Estimating hedgerow length and pattern characteristics in Great Britain using Countryside Survey data. *Journal of Environmental Management*, 60: 23–32.
- BARR, C.J., BRITT, C.P., SPARKS, T.H. & CHURCHWARD, J.M. 2005. Hedgerow Management and Wildlife: A review of research on the effects of hedgerow management and adjacent land on biodiversity. *Contract report to Defra*. 113pp.
- BAS, Y., RENARD, M., & JIGUET, F. 2009. Nesting strategy predicts farmland bird response to agricultural intensity. *Agriculture, Ecosystems and Environment*, 134: 143-147.
- BATÁRY, P., MATTIESSEN, T., & TSCHARNTKE, T. 2010. Landscape-moderated importance of hedges in conserving farmland bird diversity of organic vs. conventional croplands and grasslands. *Biological Conservation*, 143: 2020-2027.
- BAUDRY, J., BUREL, F., THENAIL, C., & LE CŒUR, D. 2000. A holistic landscape ecological study of the interactions between farming activities and ecological patterns in Brittany, France. *Landscape and Urban Planning*, 50: 119-128.
- BEALEY, C., LEDDER, E., ROBERTSON, H. & WOLTON, R. 2009. Hedgerows - their wildlife, current state and management needs. *British Wildlife*, 20: 323-329.
- BENJAMIN, R., CÉDRIC, G., & PABLO, I. 2008. Modeling spatially explicit population dynamics of *Pterostichus melanarius* I11. (Coleoptera: Carabidae) in response to changes in the composition and configuration of agricultural landscapes. *Landscape and Urban Planning*, 84: 191-199.
- BENTON, T.G., BRYANT, D.M., COLE, L. & CRICK, H.Q.P. 2012. Linking agricultural practice to insect and bird populations: a historical study over three decades. *Journal of Applied Ecology*, 39: 673–687.
- BINDING, A.E., BINDING, A., PENDLETON, T., PENDLETON, D. & CLIFTON, S.J. 2011. *The status of the hazel pot beetle Cryptocephalus coryli at Sherwood Forest National Nature Reserve*. Natural England Commissioned Reports, Number 066. 26pp.



- BISHTON, G., 1986. The diet and foraging behaviour of the dunnock *Prunella modularis* in a hedgerow habitat. *Ibis*, 128: 526-539.
- BLACKSHAW, R. P., & D'ARCY-BURT, S. 1997. Spatial distribution of bibionid larvae in agricultural grassland. *Entomologia Experimentalis Et Applicata*, 84: 17-25.
- BLAKE, D., HUTSON, A. M., RACEY, P. A., RYDELL, J., & SPEAKMAN, J. R. 1994. Use of lamplit roads by foraging bats in southern England. *Journal of Zoology*, 234: 453-462.
- BLOEMMEN M. & VAN DER SLUIS T. (Eds.) 2004. *European corridors: Strategies for corridor development for target species*. ECNC, Tilburg, the Netherlands & Alterra. 32pp.
- BOUGHEY , K.L., LAKE, I.R., HAYSON, K.A. & DOLMAN, P.M. 2011. Improving the biodiversity benefits of hedgerows: How physical characteristics and the proximity of foraging habitat affect the use of linear features by bats. *Biological Conservation*, 144: 1790–1798.
- BOUTIN, C., BARIL, A., MCCABE, S. K., MARTIN, P. A., & GUY, M. 2011. The value of woody hedgerows for moth diversity on organic and conventional farms. *Environmental Entomology*, 40: 560-569.
- BOWDEN, J.W. & DEAN, G.J.W., 1977. The distribution of flying insects in and near a tall hedgerow. *Journal of Applied Ecology*, 14: 343-354.
- BRADBURY, R. B., KYRKOS, A., MORRIS, A. J., CLARK, S. C., PERKINS, A. J., & WILSON, J. D. 2000. Habitat associations and breeding success of yellowhammers on lowland farmland. *Journal of Applied Ecology*, 37: 789-805.
- BRANDT, G., BLOWS, L., LINTON, D., PALING, N. & PRESCOTT, C. 2007. Habitat associations of British bat species on lowland farmland within the Upper Thames catchment area. *Habitat Associations of British bats. Centre for Wildlife Assessment & Conservation E-Journal* (2007), 1: 10-19.
- BRIGHT, P. & MACPHERSON, D. 2002. Hedgerow management, dormice and biodiversity. *English Nature Research Reports No 454*. English Nature, Peterborough. 33pp.
- BRIGHT, P. W. 1998. Behaviour of specialist species in habitat corridors: Arboreal dormice avoid corridor gaps. *Animal Behaviour*, 56: 1485-1490.
- BRODT, S., KLONSKY, K., JACKSON, L., BRUSH, S.B. & SMUKLER, S. 2009. Factors affecting adoption of hedgerows and other biodiversity-enhancing features on farms in California, USA. *Agroforest Syst*, 76: 195–206.
- BROOKS, D. R., BATER, J. E., CLARK, S. J., MONTEITH, D. T., ANDREWS, C., CORBETT, S. J., BEAUMONT, D.A. & CHAPMAN, J. W. 2012. Large carabid beetle declines in a United Kingdom monitoring network increases evidence for a widespread loss in insect biodiversity. *Journal of Applied Ecology*, 49: 1009-1019.
- BROWNE, S. J., AEBISCHER, N. J., YFANTIS, G., & MARCHANT, J. H. 2004. Habitat availability and use by turtle doves *Streptopelia turtur* between 1965 and 1995: An analysis of common birds census data. *Bird Study*, 51: 1-11.
- BUMBLEBEE CONSERVATION TRUST, Undated. Managing hedges and edges... for bumblebees. Factsheet, Bumblebee Conservation Trust.
http://bumblebeeconservation.org/images/uploads/Resources/BBCT_Land_Factsheet_6_Managing_hedges_edges.pdf (Accessed 18 February 2013).
- BUREL, F. & BAUDRY, J. 1990. Hedgerow networks as habitats for forest species: implications for colonising abandoned agricultural land. In: Bunce, R. & Howard, D. (Eds.), *Species Dispersal in Agricultural Habitats* pp. 238-255. Bellhaven Press, London.
- BUREL, F., & BAUDRY, J. 1990. Structural dynamic of a hedgerow network landscape in Brittany France. *Landscape Ecology*, 4: 197-210.
- BUREL, F. 1989. Landscape structure effects on carabid beetles spatial patterns in western France. *Landscape Ecology*, 2: 215-226.



- BUREL, F. 1992. Effect of landscape structure and dynamics on species diversity in hedgerow networks. *Landscape Ecology*, 6: 161-174.
- BUREL, F. 1996. Hedgerows and their role in agricultural landscapes. *Critical Reviews in Plant Sciences*, 15: 169-190.
- BUREL, F., BAUDRY, J., BUTET, A., CLERGEAU, P., DELETTRE, Y., LE COEUR, D., DUBS, F., MORVAN, N., PAILLAT, G., PETIT, S., THENAIL, C., BRUNEL, R. & LEFEUVRE, J. 1998. Comparative biodiversity along a gradient of agricultural landscapes. *Acta Oecologica*, 19: 47-60.
- BUREL, F., BUTET, A., DELETTRE, Y. R., & DE LA PEÑA, N. M. 2004. Differential response of selected taxa to landscape context and agricultural intensification. *Landscape and Urban Planning*, 67: 195-204.
- BUTET, A., PAILLAT, G. & DELETTRE, Y. 2006. Seasonal changes in small mammal assemblages from field boundaries in an agricultural landscape of western France. *Agriculture, Ecosystems & Environment*, 113: 364-369.
- CABRAL, J. A., ROCHA, A., SANTOS, M., & CRESPI, A. L. 2007. A stochastic dynamic methodology (SDM) to facilitate handling simple passerine indicators in the scope of the agri-environmental measures problematics. *Ecological Indicators*, 7: 34-47.
- CAMPAGNE, P., ROCHE, P. & TATONI, T. 2006. Factors explaining shrub species distribution in hedgerows of a mountain landscape. *Agriculture, Ecosystems & Environment*, 116: 244-250.
- CARROLL, S. 2013. *Dormice in gardens in Devon*. Interim report for Devon Mammal Group.
<http://www.devonmammalgroup.org/wp-content/uploads/DMG-grant-report-dormice-in-gardens.pdf>
- CARTER, S. P. 1990. Habitat change and bird populations. *British Wildlife*, 1: 324-335.
- CHAMBERLAIN, D. E., WILSON, J. D., & FULLER, R. J. 1999. A comparison of bird populations on organic and conventional farm systems in southern Britain. *Biological Conservation*, 88: 307-320.
- CHARRIER, S., PETIT, S., & BUREL, F. 1997. Movements of *Abax parallelepipedus* (Coleoptera, Carabidae) in woody habitats of a hedgerow network landscape: A radio-tracing study. *Agriculture, Ecosystems and Environment*, 61: 133-144.
- CHERRILL, A., MERCER, C., MCCLEAN, C., & TUDOR, G. 2001. Assessing the floristic diversity of hedge networks: A landscape perspective. *Landscape Research*, 26: 55-64.
- CHRÉTIENNE, M., & ERAUD, C. 2002. Relationship between autumn bird populations and the spatio-temporal availability of fleshy fruit in the "bocage". [Relation entre les populations d'oiseaux à l'automne-hiver et la disponibilité spatio-temporelle des fruits charnus en bocage]. *Alauda*, 70: 149-160.
- CRITCHLEY, C.N.R., WILSON, L.A., MOLE, A.C., NORTON, L.R. & SMART, S.M., 2013. A functional classification of herbaceous hedgerow vegetation for setting restoration objectives. *Biodiversity Conservation*, 22:701-717.
- CHURCHFIELD, S., HOLLIER, J., & BROWN, V K. 1997. Community structure and habitat use of small mammals in grasslands of different successional age. *Journal of Zoology*, 242: 519-530.
- CLEMENTS, D. K., & TOFTS, R. J. 1992. Hedges make the grade: A look at the wildlife value of hedges. *British Wildlife*, 4: 87-95.
- CLEMENTS, D.K. & ALEXANDER, K.N.A. 2004. *The Shapwick Project: a comparative study of the invertebrate faunas of hedgerows of different ages*. English Heritage Report, UK.
- CLEMENTS, D.K. & ALEXANDER, K.N.A. 2009. A comparative study of the invertebrate faunas of hedgerows of differing ages, with particular reference to indicators of ancient woodland and 'old growth'. *The Journal of Practical Ecology and Conservation Volume*, 8: 7-27.
- CORBETT, S. & MOLE, A. 2005. Invertebrates and hedgerows. In: Barr, C.J., Britt, C.P., Sparks, T.H. & Churchward, J.M. (Eds.), *Hedgerow management and Wildlife* 64. Defra, London. pp. 35-46.



- CORNULIER, T., ROBINSON, R.A., ELSTON, D., LAMBIN, X., SUTHERLAND, W.J. & BENTON5, T.G. 2011. Bayesian reconstitution of environmental change from disparate historical records: hedgerow loss and farmland bird declines. *Methods in Ecology and Evolution*, 2, 86–94.
- COULTHARD, E.J. 2012. Do moths use hedgerows as flight paths? An investigation of the use of linear boundary features by macro moths in intensive agricultural landscapes. In Dover, J.W. (ed.) *Hedgerow Futures: Proceedings of the first International Hedgelink Conference*, September 2012, Stoke-on-Trent, pp. 91-97.
- COWAN, A. & CROMPTON, R.M. 2004. Bats in the United Kingdom, a landscape scale perspective: considering the importance of habitat connectivity and the threats posed by fragmentation. In: Smithers, R. (ed). *Landscape Ecology of Trees and Forests: proceedings of the twelfth annual IALE (UK) conference*, held at the Royal Agricultural College, Cirencester, 21st-24th June 2004 pp. 301-306.
- CRANMER, L., MCCOLLIN, D. & OLLERTON, J. 2012. Landscape structure influences pollinator movements and directly affects plant reproductive success. *Oikos*, 121: 562-568.
- CRITCHLEY, C.N.R., WILSON, L.A., MOLE, A.C., NORTON, L.R. & SMART, S.M. 2013. A functional classification of herbaceous hedgerow vegetation for setting restoration objectives. *Biodiversity Conservation* DOI 10.1007/s10531-013-0440-5.
- CROXTON, P. J., CARVELL, C., MOUNTFORD, J. O., & SPARKS, T. H. 2002. A comparison of green lanes and field margins as bumblebee habitat in an arable landscape. *Biological Conservation*, 107: 365-374.
- CROXTON, P. J., FRANSSEN, W., MYHILL, D. G., & SPARKS, T. H. 2004. The restoration of neglected hedges: A comparison of management treatments. *Biological Conservation*, 117: 19-23.
- CROXTON, P. J., HANN, J. P., GREATOREX-DAVIES, J. N., & SPARKS, T. H. 2005. Linear hotspots? the floral and butterfly diversity of green lanes. *Biological Conservation*, 121: 579-584.
- DANIELS, R. J. R. 1994. A landscape approach to conservation of birds. *Journal of Biosciences*, 19: 503-509.
- DAVEY, C. M., VICKERY, J. A., BOATMAN, N. D., CHAMBERLAIN, D. E., & SIRIWARDENA, G. M. 2010. Entry level stewardship may enhance bird numbers in boundary habitats. *Bird Study*, 57:415-420.
- DAVEY, C., VICKERY, J., BOATMAN, N., CHAMBERLAIN, D., PARRY, H. & SIRIWARDENA, G. 2010. Regional variation in the efficacy of Entry Level Stewardship in England. *Agriculture, Ecosystems and Environment*, 139: 121–128.
- DAVEY, C.M., VICKERY, J.A., BOATMAN, N.D., CHAMBERLAIN, D.E., PARRY, H.R. & SIRIWARDENA, G.M. 2010. Assessing the impact of Entry Level Stewardship on lowland farmland birds in England. *Ibis*, 152: 459–474.
- DAVIES, Z. & PULLIN, A. 2007. Are hedgerows effective corridors between fragments of woodland habitat? An evidence-based approach. *Landscape Ecology*, 22: 333-351.
- DAVIES, Z.G. & PULLIN, A.S. 2006. Do Habitat Corridors Increase Population Viability? Part A: Do Hedgerow Corridors Increase The Population Viability Of Woodland Species? CEE review 05-001 (SR8a). Collaboration for Environmental Evidence: <http://www.environmentalevidence.org/Documents/Summary/Summary-SR8a.pdf> (Accessed 18 February 2013).
- DAVIES, Z.G., TYLER, C., STEWART, G.B. & PULLIN, A.S. 2006. Are current management recommendations for conserving saproxylic invertebrates effective? CEE review 05-011 (SR17). *Collaboration for Environmental Evidence*: www.environmentalevidence.org/SR17.html.
- DAWSON, D. 1994. Are habitat corridors conduits for animals and plants in a fragmented landscape? A review of the scientific evidence. *English Nature Research Report* 94. 89pp.
- DE VRIES, H. H., ENS, S. H., DE GRAAF, G., TEUNISSEN, L., VAN DER VELDE, R., VOGELAAR, L., WINTERINK, A. & VISSER, M. E. 2011. Synchronisation of egg hatching of brown hairstreak (*Thecla betulae*) and budburst of blackthorn (*Prunus spinosa*) in a warmer future. *Journal of Insect Conservation*, 15: 311-319.



- DECKERS, B., HERMY, M. & MUYS, B. 2004. Factors affecting plant species composition of hedgerows: relative importance and hierarchy. *Acta Oecologica*, 26, 23-37.
- DECKERS, B., KERSELAERS, E., GULINCK, H., MUYS, B., & HERMY, M. 2005. Long-term spatio-temporal dynamics of a hedgerow network landscape in Flanders, Belgium. *Environmental Conservation*, 32: 20-29.
- DECKERS, B., VERHEYEN, K., HERMY, M. & MUYS, B. 2004. Differential environmental response of plant functional types in hedgerow habitats. *Basic and Applied Ecology*, 5: 551—566.
- DEFRA 2012. A Strategy for England's wildlife and ecosystem services. Biodiversity 2020 Indicators: 2012 Assessment. Defra, London 142pp. <http://www.Defra.gov.uk/statistics/environment/biodiversity/england-biodiversity-indicators/>
- DEFRA 2007. *Hedgerow Survey Handbook: a standard procedure for local surveys in the UK*. Defra, London. 143pp <http://archive.defra.gov.uk/foodfarm/landmanage/landscape/documents/hedgerow-survey-handbook.pdf> (Accessed 18 February 2012).
- DELETTRE, Y. R., & MORVAN, N. 2000. Dispersal of adult aquatic Chironomidae (Diptera) in agricultural landscapes. *Freshwater Biology*, 44:399-411.
- DELETTRE, Y. R. 2005. Short-range spatial patterning of terrestrial Chironomidae (Insecta: Diptera) and farmland heterogeneity. *Pedobiologia*, 49: 15-27.
- DICKS, L.V., ABRAHAMS, A., ATKINSON, J., BIESMEIJER, J., BOURN, N., BROWN, C., BROWN, M.J.F., CARVELL, C., CONNOLLY, C., CRESSWELL, J.E., CROFT, P., DARVILL, B., ZYLVIA, P., EFFINGHAM, P., FOUNTAIN, M., GOGIN, A., HARDING, D., HARDING, T., HARTFIELD, C., HEARD, M.S., HEATHCOTE, R., HEAVER, D., HOLLAND, J., HOWE, M., HUGHES, B., HUXLEY, T., KUNIN, W.E., LITTLE, J., MASON, C., MEMMOTT, J., OSBORNE, J., PANKHURST, T., PAXTON, R.J., POCOCK, M.J.O., POTTS, S.G., POWER, E.F., RAINES, N.E., RANELAGH, E., ROBERTS, S., SAUNDERS, R., SMITH, K., SMITH, R.M., SUTTON, P., TILLEY, L.A.N., TINSLEY, A., TONHASCA, A., VANBERGEN, A.J., WEBSTER, S., WILSON, A. & SUTHERLAND, W.J. 2012. Identifying key knowledge needs for evidence-based conservation of wild insect pollinators: a collaborative cross-sectoral exercise. *Insect Conservation and Diversity* doi: 10.1111/j.1752-4598.2012.00221.x.
- DOUGLAS, D.J.T., MOREBY, S.J. & BENTON, T.G. 2011. Provisioning with cereal grain depresses the body condition of insectivorous Yellowhammer *Emberiza citrinella* nestlings. *Bird Study*, 59: 105-109.
- DOUGLAS, D.J.T., VICKERY, J.A. & BENTON, T.G. 2009. Improving the value of field margins as foraging habitat for farmland birds. *Journal of Applied Ecology*, 46: 353–36.
- DOVER, J.W. 1990. Butterflies and wildlife corridors. In: NODDER, C. (ed). The Game Conservancy Review of 1989. The Game Conservancy, Fordingbridge, pp 62-64.
- DOVER, J. 1996. Factors affecting the distribution of butterflies on arable farmland. *Journal of Applied Ecology* 54: 117-124.
- DOVER, J. W. 1997. Conservation headlands: Effects on butterfly distribution and behaviour. *Agriculture, Ecosystems and Environment*, 63: 31-49.
- DOVER, J., SOTHERTON, N. & GOBBETT, K. 1990. Reduced pesticide inputs on cereal field margins: the effects on butterfly abundance. *Ecological Entomology*, 15:17-24.
- DOVER, J. W., SPARKS, T. H., & GREATOREX-DAVIES, J. N. 1997. The importance of shelter for butterflies in open landscapes. *Journal of Insect Conservation*, 1: 89-97.
- DOVER, J., & SPARKS, T. 2000. A review of the ecology of butterflies in British hedgerows. *Journal of Environmental Management*, 60: 51-63.
- DOVER, J., SPARKS, T., CLARKE, S., GOBBETT, K., & GLOSSOP, S. 2000. Linear features and butterflies: The importance of green lanes. *Agriculture, Ecosystems and Environment*, 80: 227-242.
- DOVER, J. W., & FRY, G. L. A. 2001. Experimental simulation of some visual and physical components of a hedge and the effects on butterfly behaviour in an agricultural landscape. *Entomologia Experimentalis Et Applicata*, 100: 221-233.



- DOVER, J.W. & SPARKS, T.H. 2001. Green lanes: biodiversity reservoirs in farmland? In: BARR, C.J. & PETIT, S. (eds) *Hedgerows of the World: their ecological functions in different landscapes*. IALE(UK), Lymm, pp 241-250.
- DOVER, J. & SETTELE, J. 2009. The influences of landscape structure on butterfly distribution and movement: a review. *Journal of Insect Conservation*, 13:3-27.
- DOWNS, N. C., & RACEY, P. A. 2006. The use by bats of habitat features in mixed farmland in Scotland. *Acta Chiropterologica*, 8(1), 169-185.
- DRAKE, C.M. 2001. The importance of temporary waters for Diptera (True-Flies). *Freshwater Forum*, 17: 26-39.
- DUBOIS, G. F., VIGNON, V., DELETTRE, Y. R., RANTIER, Y., VERNON, P., & BUREL, F. 2009. Factors affecting the occurrence of the endangered saproxylic beetle *Osmoderma eremita* (Scopoli, 1763) (Coleoptera: Cetoniidae) in an agricultural landscape. *Landscape and Urban Planning*, 91: 152-159.
- DUELLI, P. 1990. Population movements of arthropods between natural and cultivated areas. *Biological Conservation*, 54: 193-207.
- DUNN, J. C., & MORRIS, A. J. 2012. Which features of UK farmland are important in retaining territories of the rapidly declining turtle dove *Streptopelia turtur*? *Bird Study*, 59: 394-402.
- DUNN, J.C., HAMER, K.C. & BENTON, T.G. 2010. Fear for the family has negative consequences: indirect effects of nest predators on chick growth in a farmland bird. *Journal of Applied Ecology*, 47: 994–1002.
- EBERT, J., & MÜLLER-PFANNENSTIEL, K. 2008. Transfer of tree roots colonised with larvae of the stag beetle - A measure to reduce damage for a species protected according to the Habitats Directive. [Umsetzung von mit Hirschläufer-Larven Besetzten Baumwurzeln - Eine Maßnahme zur Schadensbegrenzung für eine FFH-Art]. *Naturschutz Und Landschaftsplanung*, 40(4), 106-112.
- EDGAR, P., FOSTER, J. & BAKER, J. 2010. *Reptile Habitat Management Handbook*. Amphibian and Reptile Conservation, Bournemouth, 84pp.
- EGGERS, B., MATERN, A., DREES, C., EGGERS, J., HÄRDLE, W., & ASSMANN, T., 2010. Value of semi-open corridors for simultaneously connecting open and wooded habitats: A case study with ground beetles: Contributed paper. *Conservation Biology*, 24: 256-266.
- EHLERS, S. 2012. The importance of hedgerows for hazel dormice (*Muscardinus avellanarius*) in Northern Germany. *Peckiana*, 8: 41-47.
- ENTWISTLE, A.C., RACEY, P.A. & SPEAKMAN, J.R. 1996. Habitat exploitation by a gleaning bat, *Plecotus auritus*. *Philosophical Transactions of the Royal Society B: Biological Sciences*, 351: 921-931.
- EVANS, D.M., POCOCK, M.J.O, BROOKS, J. & MEMMOTT, J. 2011. Seeds in farmland food-webs: Resource importance, distribution and the impacts of farm management. *Biological Conservation*, 144: 2941–2950.
- EVANS, D.M., POCOCK, M.J.O, BROOKS, J. & MEMMOTT, J. In press. The robustness of a network of ecological networks to habitat loss.
- EYBERT, M.C., CONSTANT, P., & LEFEUVRE, J.C. 1995. Effects of Changes in Agricultural Landscape on a Breeding Population of Linnets *Acanthis-Cannabina* L. Living in Adjacent Heathland. *Biological Conservation*, 74: 195-202.
- FARTMANN, T., & TIMMERMANN, K. 2006. Where to find the eggs and how to manage the breeding sites of the brown hairstreak (*Thecla betulae* (Linnaeus, 1758)) in central Europe? *Nota Lepidopterologica*, 29: 125-134.
- FEBER, R. E., JOHNSON, P. J., FIRBANK, L. G., HOPKINS, A., & MACDONALD, D. W. 2007. A comparison of butterfly populations on organically and conventionally managed farmland. *Journal of Zoology*, 273: 30-39.
- FIELD, R. G., & MASON, C. F. 2005. The utilization of two-metre countryside stewardship scheme grass margins by the gatekeeper *Pyronia tithonus* (L.). *Journal of Natural History*, 39: 1533-1538.



- FIELD, R. G., GARDINER, T., MASON, C. F., & HILL, J. 2006. Countryside stewardship scheme and butterflies: A study of plant and butterfly species richness. *Biodiversity and Conservation*, 15: 443-452.
- FIRBANK, L. G., BARR, C. J., BUNCE, R. G. H., FURSE, M. T., HAINES-YOUNG, R., HORNUNG, M., HOWARD, D.C., SHEAIL, J., SIER, A. & SMART, S. M. 2003. Assessing stock and change in land cover and biodiversity in GB: An introduction to countryside survey 2000. *Journal of Environmental Management*, 67: 207-218.
- FOREST RESEARCH 2009. Trends, long term survival and ecological values of hedgerow trees: development of populations models to inform strategy. *Defra research project BD211*.
<http://www.hedgelink.org.uk/files/Hedgrow%20tree%20BD2111%20sid5%20final%20report1,%20Forest%20Research,%20Aug%2010.pdf>
- FORMAN, R.T.T. & BAUDRY, J. 1984. Hedgerows and hedgerow networks in landscape ecology. *Environmental Management*, 8: 495-510.
- FOURNIER, E., & LOREAU, M. 1999. Effects of newly planted hedges on ground-beetle diversity (Coleoptera, Carabidae) in an agricultural landscape. *Ecography*, 22: 87-97.
- FOURNIER, E. & LOREAU, M. 2001. Respective roles of recent hedges and forest patch remnants in the maintenance of ground-beetle (Coleoptera: Carabidae) diversity in an agricultural landscape. *Landscape Ecology*, 16: 17-32.
- FREMLIN, M. & HENDRIKS, P. 2011. Sugaring for stag beetles – different feeding strategies of *Lucanus cervus* and *Dorcus parallelipipedus*. *Bulletin of the Amateur Entomologists' Society*, 70: 57-67.
- FROUZ, J. & PAOLETTI, M.G. 2000. Spatial distribution of different life stages of one Dipteron community along hedgerow and field margin. *Landscape and Urban Planning*, 48: 19-29.
- FRY, R. & LONSDALE, D. (Eds.), 1991. *Habitat Conservation for Insects- a neglected green issue*. Amateur Entomologist Society, Middlesex.
- FUENTES-MONTEMAYOR, E., GOULSON, D. & PARK, K.J. 2011. Pipistrelle bats and their prey do not benefit from four widely applied agri-environment management prescriptions. *Biological Conservation*, 144: 2233–2246.
- FUENTES-MONTEMAYOR, E., GOULSON, D., & PARK, K. J. 2011. The effectiveness of agri-environment schemes for the conservation of farmland moths: Assessing the importance of a landscape-scale management approach. *Journal of Applied Ecology*, 48: 532-542.
- FULLER, R. J., CHAMBERLAIN, D. E., BURTON, N. H. K., & GOUGH, S. J. 2001. Distributions of birds in lowland agricultural landscapes of England and Wales: How distinctive are bird communities of hedgerows and woodland? *Agriculture, Ecosystems and Environment*, 84: 79-92.
- FULLER, R. J., TREVELYAN, R. J., & HUDSON, R. W. 1997. Landscape composition models for breeding bird populations in lowland English farmland over a 20 year period. *Ecography*, 20: 295-307.
- GABRIEL, D., SAIT, S.M., HODGSON, J.A., SCHMUTZ, U., KUNIN, W.E. & BENTON, T.G. 2010. Scale matters: the impact of organic farming on biodiversity at different spatial scales. *Ecology Letters*, 13: 858–869.
- GARCÍA, A. F., GRIFFITHS, G. J. K., & George Thomas, C. F. 2000. Density, distribution and dispersal of the carabid beetle *Nebria brevicollis* in two adjacent cereal fields. *Annals of Applied Biology*, 137: 89-97.
- GELLING, M., MACDONALD, D.W. & MATHEWS, F., 2007. Are hedgerows the route to increased farmland small mammal density? Use of hedgerows in British pastoral habitats. *Landscape Ecology*, 22: 1019–1032.
- GILLINGS, S., & FULLER, R. J. 1998. Changes in bird populations on sample lowland English farms in relation to loss of hedgerows and other non-crop habitats. *Oecologia*, 116: 120-127.
- GILROY, J. J., ANDERSON, G. Q. A., GRICE, P. V., VICKERY, J. A., & SUTHERLAND, W. J. 2010. Mid-season shifts in the habitat associations of yellow wagtails *Motacilla flava* breeding in arable farmland. *Ibis*, 152: 90-104.
- GOITI, U., AIHARTZA, J. R., GARIN, I., & ZABALA, J. 2003. Influence of habitat on the foraging behavior of the Mediterranean horseshoe bat, *Rhinolophus euryale*. *Acta Chiropterologica*, 5: 75-84.



- GOITI, U., GARIN, I., ALMENAR, D., SALSAMENDI, E., & AIHARTZA, J. 2008. Foraging by Mediterranean horseshoe bats (*Rhinolophus euryale*) in relation to prey distribution and edge habitat. *Journal of Mammalogy*, 89: 493-502.
- GREEN, R. E., OSBORNE, P. E., & SEARS, E. J. 1994. The distribution of passerine birds in hedgerows during the breeding season in relation to characteristics of the hedgerow and adjacent farmland. *Journal of Applied Ecology*, 31: 677-692.
- GRIFFITHS, G. J. K., WINDER, L., HOLLAND, J. M., THOMAS, C. F. G., & WILLIAMS, E. 2007. The representation and functional composition of carabid and staphylinid beetles in different field boundary types at a farm-scale. *Biological Conservation*, 135: 145-152.
- GRIXTI, J. C., WONG, L. T., CAMERON, S. A., & FAVRET, C. 2009. Decline of bumble bees (*Bombus*) in the North American Midwest. *Biological Conservation*, 142:75-84.
- GROOT, J.C.J., JELLEMA, A. & ROSSING, W.A.H. 2008. Designing a hedgerow network in a multifunctional agricultural landscape: Balancing trade-offs among ecological quality, landscape character and implementation costs. *European Journal of Agronomy*, 32: 112–119.
- HAALAND, C. & BERSIER, L.F. 2011. What can sown wildflower strips contribute to butterfly conservation?: an example from a Swiss lowland agricultural landscape. *Journal of Insect Conservation*, 15:301–309.
- HEARD, M.S., BOTHAM, M, BROUGHTON, R., CARVELL, C., HINSLEY, S., WOODCOCK, B., PYWELL, R.F. 2012. *Quantifying the effects of Entry Level Stewardship (ELS) on biodiversity at the farm scale: the Hillesden Experiment*. NERC Centre for Ecology and Hydrology. Report to Defra and Natural England.
- HANLEY, M. E., FRANCO, M., DEAN, C. E., FRANKLIN, E. L., HARRIS, H. R., HAYNES, A. G., RAPSON, S. R., ROWSE, G., THOMAS, K. C., WATERHOUSE, B. R. & KNIGHT, M. E. 2011. Increased bumblebee abundance along the margins of a mass flowering crop: Evidence for pollinator spill-over. *Oikos*, 120: 1618-1624.
- HANNON, L. E., & SISK, T. D. 2009. Hedgerows in an agri-natural landscape: Potential habitat value for native bees. *Biological Conservation*, 142: 2140-2154.
- HARRIS, G. L., ROSE, S. C., PARISH, T., & MOUNTFORD, J. O. 1991. A case study observing changes in land drainage and management in relation to ecology. *Hydrological Basis of Ecologically Sound Management of Soil and groundwater. Proceedings of Symposium, Vienna, 1991*. pp 247-256.
- HARVEY, D. J., & GANGE, A. C. 2006. Size variation and mating success in the stag beetle, *Lucanus cervus*. *Physiological Entomology*, 31: 218-226.
- HARVEY, D. J., GANGE, A. C., HAWES, C. J., & RINK, M. 2011. Bionomics and distribution of the stag beetle, *Lucanus cervus* (L.) across Europe. *Insect Conservation and Diversity*, 4: 23-38.
- HARVEY, D. J., HAWES, C. J., GANGE, A. C., FINCH, P., CHESMORE, D., & FARR, I. 2011. Development of non-invasive monitoring methods for larvae and adults of the stag beetle, *Lucanus cervus*. *Insect Conservation and Diversity*, 4: 4-14.
- HARVEY, D.J. & GANGE, A.C. 2011. The stag beetle: a collaborative conservation study across Europe. *Insect Conservation and Diversity*, 4, 2–3
- HARVEY, D.J., GANGE, A.C., HAWES, C.J. & RINK, M. 2011. Bionomics and distribution of the stag beetle, *Lucanus cervus* (L.) across Europe. *Insect Conservation and Diversity*, 4: 23–38.
- HAWES, C. J. 2008. The stag beetle *Lucanus cervus* (Linnaeus, 1758) (Coleoptera: Lucanidae): A mark-release-recapture study undertaken in one United Kingdom residential garden. *Revue d'Ecologie (La Terre Et La Vie)*, 63: 139-146.
- HEINEIN, K., WEGNER, J. & MERRIAM, G. 1998. Population effects of landscape model manipulation on two behaviourally different woodland small mammals. *Oikos*, 81: 168 – 186.
- HINSLEY, S. A. & BELLAMY, P. E. 2000. The influence of hedge structure, management and landscape context on the value of hedgerows to birds: A review. *Journal of Environmental Management*, 60: 33–49.
- HODGSON, J.A., KUNIN, W.E., THOMAS, C.D., BENTON, T.G. & GABRIEL, D. 2010. Comparing organic farming and land sparing: optimizing yield and butterfly populations at a landscape scale. *Ecology Letters*, 13: 1358-67.



HOF, A.R., 2009. A study of the current status of the hedgehog (*Erinaceus europaeus*), and its decline in Great Britain since 1960. *Thesis submitted for the degree of Doctor of Philosophy* (PhD) Royal Holloway, University of London Egham, Surrey, TW20 0EX, UK

HOF, A.R., AND BRIGHT, P.W. 2010. The value of agri-environment schemes for macroinvertebrate feeders: hedgehogs on arable farms in Britain. *Animal Conservation*, 13:467-473.

HOLDERIED, M. W., JONES, G., & VON HELVERSEN, O. 2006. Flight and echolocation behaviour of whiskered bats commuting along a hedgerow: Range-dependent sonar signal design, doppler tolerance and evidence for 'acoustic focusing'. *Journal of Experimental Biology*, 209: 1816-1826.

HOLLAND, J.M., BEGBIE, M., BIRKETT, T., REYNOLDS, C.J.M. & THOMAS, C.F.G. 2001. The influence of hedgerows on coleopteran distributions: results from a multi-field sampling study. The Game Conservancy Trust & Seale-Hayne Faculty, University of Plymouth. 12pp. Publishing source uncertain.

http://www.rothamsted.ac.uk/pie/sadie/reprints/holland_hedgerows_2001.pdf

HOLLAND, J.M., OATEN, H, MOREBY, S., BIRKETT, T., SIMPER, J., SOUTHWAY, S. & SMITH, B.M. 2012. Agri-environment scheme enhancing ecosystem services: A demonstration of improved biological control in cereal crops. *Agriculture, Ecosystems and Environment*, 155: 147–152.

HOLZSCHUH, A., STEFFAN-DEWENTER, I., & TSCHARNTKE, T. 2009. Grass strip corridors in agricultural landscapes enhance nest-site colonization by solitary wasps. *Ecological Applications*, 19:123–132.

HOPWOOD, J. L. 2008. The contribution of roadside grassland restorations to native bee conservation. *Biological Conservation*, 141: 2632-2640.

HRADETZKY, R & KROMP, B. 1997. Spatial distribution of flying insects in an organic rye field and an adjacent hedge and forest edge. *Entomological Research in Organic Agriculture*, 1997: 353-357.

HULL, S.K. 1991. A study undertaken in southern Britain to investigate the late-flushing of hedgerow ash in spring 1987. *Forestry*, 64: 189-197.

JACOBS, J. H., CLARK, S. J., DENHOLM, I., GOULSON, D., STOATE, C., & OSBORNE, J. L. 2010. Pollinator effectiveness and fruit set in common ivy, *Hedera helix* (Araliaceae). *Arthropod-Plant Interactions*, 4: 19-28.

JACOBS, J.H., CLARK, S.J., DENHOLM, I., GOULSON, D., STOATE, C. & OSBORNE, J.L. 2009. Pollination biology of fruit-bearing hedgerow plants and the role of flower-visiting insects in fruit-set. *Annals of Botany*, 104: 1397–1404.

JOBIN, B., CHOINIÈRE, L., & BÉLANGER, L. 2001. Bird use of three types of field margins in relation to intensive agriculture in Quebec, Canada. *Agriculture, Ecosystems and Environment*, 84: 131-143.

JOFRÉ, G.M. & Reading, C.J. 2012. An assessment of the impact of conservation grazing on reptile populations. *Amphibian and Reptile Conservation Research Report 12/01*. 48pp.

JOLY, P., MIAUD, C., LEHMANN, A. & GROLET, O. 2001. Habitat matrix effects on pond occupancy in newts. *Conservation Biology*, 15: 239-248.

JOPP, F., & REUTER, H. 2005. Dispersal of carabid beetles - emergence of distribution patterns. *Ecological Modelling*, 186: 389-405.

JOYCE, K. A., HOLLAND, J. M., & DONCASTER, C. P. 1999. Influences of hedgerow intersections and gaps on the movement of Carabid beetles. *Bulletin of Entomological Research*, 89: 523-531.

JOYCE, K.A. 1997. The role of hedgerows in the ecology of invertebrates in arable landscapes. Unpublished PhD study, University of Southampton.

JOYCE, K.A., JEPSON, P.C., DONCASTER, C.P. & HOLLAND, J.M.. 1997. Arthropod distribution patterns and dispersal processes within the hedgerow. In: Cooper, A. & Power, J. (Eds.), *Species Dispersal and Land Use Processes* pp. 103-110. International Association for Land Use Ecology, Coleraine, UK.



- KELLEHER, K. M., & O'HALLORAN, J. 2007. Influence of nesting habitat on breeding song thrushes *Turdus philomelos*. *Bird Study*, 54: 221-229.
- KIRBY, K.J. & DRAKE, C.M. 1993. Dead wood matters: the ecology and conservation of saproxylic invertebrates in Britain. *English Nature Science Report No. 7*. English Nature, Peterborough. 105pp.
- KLEIJN, D., BAQUERO, R.A., CLOUGH, Y., DÍAZ, M., DE ESTEBAN, J., FERNÁNDEZ, F., GABRIEL, D., HERZOG, F., HOLZSCHUH, A., JÖHL, R., KNOP, E., KRUESS, A., MARSHALL, E.J.P., STEFFAN-DEWENTER, I., TSCHARNTKE, T., VERHULST, J., WEST, T.M. & YELA, J.L. 2006. Mixed biodiversity benefits of agri-environment schemes in five European countries. *Ecology Letters*, 9: 243-254.
- KNIGHT, T. 2006. The use of landscape features and habitats by the lesser horseshoe bat (*Rhinolophus hipposideros*). Unpubl. PhD thesis, University of Bristol.
- KOTZAGEORGIS, G. C., & Mason, C. F. 1997. Small mammal populations in relation to hedgerow structure in an arable landscape. *Journal of Zoology*, 242: 425-434.
- KREMEN, C. Undated. Pollinator conservation and our food security: what do hedgerows contribute? <http://nature.berkeley.edu/kremenlab/pdf/Hedgerow%20Native%20Bee%20Project.pdf>
- KRISTENSEN, L. S., THENAIL, C., & KRISTENSEN, S. P. 2004. Landscape changes in agrarian landscapes in the 1990s: The interaction between farmers and the farmed landscape. A case study from Jutland, Denmark. *Journal of Environmental Management*, 71: 231-244.
- KRISTENSEN, S. P. 2003. Multivariate analysis of landscape changes and farm characteristics in a study area in central Jutland, Denmark. *Ecological Modelling*, 168: 303-318.
- KYRKOS, A., WILSON, J. D., & FULLER, R. J. 1998. Farmland habitat change and abundance of yellowhammers *Emberiza citrinella*: An analysis of common birds census data. *Bird Study*, 45: 232-246.
- LACK, P. C. 1987. The effects of severe hedge cutting on a breeding bird population. *Bird Study*, 34: 139-146.
- LAWTON, J.H., BROTHERTON, P.N.M., BROWN, V.K., ELPHICK, C., FITTER, A.H., FORSHAW, J., HADDOW, R.W., HILBORNE, S., LEAFE, R.N., MACE, G.M., SOUTHGATE, M.P., SUTHERLAND, W.A., TEW, T.E., VARLEY, J., & WYNNE, G.R. 2010. *Making Space for Nature: a review of England's wildlife sites and ecological network*. Report to Defra. 119pp. <http://archive.defra.gov.uk/environment/biodiversity/documents/201009space-for-nature.pdf> (Accessed 18 February 2013).
- LE CŒUR, D., BAUDRY, J., BUREL, F., & THENAIL, C. 2002. Why and how we should study field boundary biodiversity in an agrarian landscape context. *Agriculture, Ecosystems and Environment*, 89: 23-40.
- LEWIS, T. 1969. The distribution of flying insects near a low hedgerow. *Journal of Applied Ecology*, 6: 443-452.
- LEWIS, T. 1970. Patterns of distribution of insects near a windbreak of tall trees. *Annals of Applied Biology*, 65: 213-220.
- LIMPENS, H.J.G.A. & KAPTEYN, K. 1991. Bats, their behaviour and linear landscape elements. *Myotis* 29, 63-71.
- LONGLEY, M., 2003. *Greater horseshoe bat Project*, 1998-2003. English Nature Research Report 532. 38pp.
- LUDWIG, M., SCHLINKERT, H., HOLZSCHUH, A., FISCHER, C., SCHERBER, C., TRNKA, A., TSCHARNTKE, T. & BATÁRY, P. 2012. Landscape-moderated bird nest predation in hedges and forest edges. *Acta Oecologica*, 45: 50-56.
- LUFF, M.L. & WOIWOOD, I.P. 1995. Insects as indicators of land use change: a European perspective, focusing on moths and ground beetles. In: Harrington, R. & Stork, N.E. (Eds.), *Insects in a Changing Environment*. pp. 399-422. Academic Press, London.
- LYE, G., PARK, K., OSBORNE, J., HOLLAND, J., & GOULSON, D. 2009. Assessing the value of rural stewardship schemes for providing foraging resources and nesting habitat for bumblebee queens (Hymenoptera: Apidae). *Biological Conservation*, 142: 2023-2032.



- MACDONALD, D. W., & JOHNSON, P. J. 1995. The relationship between bird distribution and the botanical and structural characteristics of hedges. *Journal of Applied Ecology*, 32: 492-505.
- MACDONALD, D.W. TATTERSALL, F.H. SERVICE, K.M. FIRBANK, L.G. & FEBER, R.E. 2007. Mammals, agri-environment schemes and set-aside – what are the putative benefits? *Mammal Review*, 37: 259–277.
- MACDONALD, D.W., TEW, T.E. & TODD, I.A. 2004. The ecology of weasels (*Mustela nivalis*) on mixed farmland in southern England. *Biologia, Bratislava*, 59: 235-241.
- MACLEOD, A. 1999. Attraction and retention of *Episyrrhus balteatus* DeGeer (Diptera: Syrphidae) at an arable field margin with rich and poor floral resources. *Agriculture, Ecosystems and Environment*, 73: 241.
- MARSHALL, E.J.R. & MOONEN, A.C. 2002. Field margins in Northern Europe: their functions and interactions with agriculture. *Agriculture, Ecosystems & Environment*, 89: 5-21.
- MARSLAND, A. 1989. Historic hedgerows in Rutland: The character and ecology of the woodland relic hedgerow. *Rutland Record*, 9: 299-304.
- MARTIN, M., BASTARDIE, F., RICHARD, D., & BUREL, F. 2001. Studying boundary effects on animal movement in heterogeneous landscapes: The case of *Abax ater* (Coleoptera: Carabidae) in hedgerow network landscapes. *Comptes Rendus De l'Academie Des Sciences - Serie III*, 324: 1029-1035.
- MASON, C. F., & MACDONALD, S. M. 2000. Influence of landscape and land-use on the distribution of breeding birds in farmland in eastern England. *Journal of Zoology*, 251: 339-348.
- MAUDSLEY, M., SEELEY, B., & LEWIS, O. 2002. Spatial distribution patterns of predatory arthropods within an English hedgerow in early winter in relation to habitat variables. *Agriculture, Ecosystems and Environment*, 89: 77-89.
- MAUDSLEY, M.J. 2000. A review of the ecology and conservation of hedgerow invertebrates in Britain. *Journal of Environmental Management*, 60: 65–76.
- MAUREMOOTO, J. R., WRATTEN, S. D., WORNER, S. P., & FRY, G. L. 1995. Permeability of hedgerows to predatory carabid beetles. *Agriculture, Ecosystems and Environment*, 52: 141-148.
- MCCOLLIN, D., JACKSON, J. I., BUNCE, R. G. H., BARR, C. J., & STUART, R. 2000. Hedgerows as habitat for woodland plants. *Journal of Environmental Management*, 60: 77-90.
- MEISTER, B., HOFER, U., URSENBACHER, S. & BAUR, B. 2010. Spatial genetic analysis of the grass snake, *Natrix natrix* (Squamata: Colubridae), in an intensively used agricultural landscape. *Biological Journal of the Linnean Society*, 101: 51–58.
- MEMMOTT, J. 1999. The structure of a plant-pollinator food web. *Ecology Letters*, 2: 276-280
- MERCKX, T. & BERWAERTS, K. 2010. What type of hedgerows do Brown hairstreak (*Thecla betulae* L.) butterflies prefer? Implications for European agricultural landscape conservation. *Insect Conservation and Diversity*, (2010) doi: 10.1111/j.1752-4598.2010.00088.x
- MERCKX, T., & VAN DYCK, H. 2005. Mate location behaviour of the butterfly *Pararge aegeria* in woodland and fragmented landscapes. *Animal Behaviour*, 70: 411-416.
- MERCKX, T., FEBER, R. E., DULIEU, R. L., TOWNSEND, M. C., PARSONS, M. S., BOURN, N. A. D., RIORDAN, P. & MACDONALD, D. W. 2009. Effect of field margins on moths depends on species mobility: Field-based evidence for landscape-scale conservation. *Agriculture, Ecosystems and Environment*, 129: 302-309.
- MERCKX, T., FEBER, R. E., PARSONS, M. S., BOURN, N. A. D., TOWNSEND, M. C., RIORDAN, P., & MACDONALD, D. W. 2010. Habitat preference and mobility of *Polia bombycina*: Are non-tailored agri-environment schemes any good for a rare and localised species? *Journal of Insect Conservation*, 14: 499-510.
- MERCKX, T., FEBER, R.E, RIORDAN, P., TOWNSEND, M.C., BOURN, N.A.D., PARSONS, M.S. & MACDONALD, D.W. 2009. Optimizing the biodiversity gain from agri-environment schemes. *Agriculture, Ecosystems and Environment*, 130: 177–182.



- MERCKX, T., FEBERA, R.E., MCLAUGHLAN, C., BOURN, N.A.D., PARSONS, M.S., TOWNSEND, M.C., RIORDAN, P. & MACDONALD, D.W. 2010. Shelter benefits less mobile moth species: The field-scale effect of hedgerow trees. *Agriculture, Ecosystems and Environment*, 138: 147–151.
- MERCKX, T., MARINI, L., FEBER, R.E. & MACDONALD, D.W. 2012. Hedgerow trees and extended-width field margins enhance macro-moth diversity: implications for management. *Journal of Applied Ecology*, 49: 1396–1404.
- MEYLING, N. V., LÜBECK, M., BUCKLEY, E. P., EILENBERG, J., & REHNER, S. A. 2009. Community composition, host range and genetic structure of the fungal entomopathogen beauveria in adjoining agricultural and seminatural habitats. *Molecular Ecology*, 18: 1282-1293.
- MILLÁN DE LA PEÑA, N., BUTET, A., DELETTRE, Y., MORANT, P., & BUREL, F. 2003. Landscape context and carabid beetles (coleoptera: Carabidae) communities of hedgerows in western France. *Agriculture, Ecosystems and Environment*, 94: 59-72.
- MORANDIN, L. A., & KREMEN, C. 2013 Bee preference for native versus exotic plants in restored agricultural hedgerows. *Restoration Ecology*, 21: 26–32.
- MORTELLITI, A., FAGIANI, S., BATTISTI, C., CAPIZZI, D., & BOITANI, L. 2010. Independent effects of habitat loss, habitat fragmentation and structural connectivity on forest-dependent birds. *Diversity and Distributions*, 16: 941-951.
- MOTTE, G., & LIBOIS, R. 2002. Conservation of the lesser horseshoe bat (*Rhinolophus hipposideros* bechstein, 1800) (Mammalia: Chiroptera) in Belgium. A case study of feeding habitat requirements. *Belgian Journal of Zoology*, 132: 49-54.
- Muir, R. & Muir, N., 1987. *Hedgerows: Their History and Wildlife*. Michael Joseph, UK.
- MURPHY, S. E., GREENAWAY, F. & HILL, D. A., 2012. Patterns of habitat use by female brown long-eared bats presage negative impacts of woodland conservation management. *Journal of Zoology*, 288: 177-183.
- NAGY, Z.T. & KORSOS, Z. 1998. Data on movements & thermal biology of grass snake (*Natrix natrix* L.) using radiotelemetry. In Miaud, C. & Guyetent, R. (eds) Le Bourget Du Lac, France, pp. 339-343.
- NEW, T.R. 2004. Moths (Insecta: Lepidoptera) and conservation: background and perspective. *Journal of Insect Conservation*, 8: 79-94.
- NEWTON, I. 2004. The recent declines of farmland bird populations in Britain: An appraisal of causal factors and conservation actions. *Ibis*, 146: 579-600.
- NORTON, L., JOHNSON, P., JOYS, A., STUART, R., CHAMBERLAIN, D., FEBER, R., FIRBANK, L., MANLEY, W., WOLFE, M., HART, B., MATHEWS, F., MACDONALD, D. & FULLER, R.J. 2009. Consequences of organic and non-organic farming practices for field, farm and landscape complexity. *Agriculture, Ecosystems and Environment*, 129: 221–227.
- OAKELEY, S. F., & JONES, G. 1998. Habitat around maternity roosts of the 55 kHz phonic type of pipistrelle bats (*Pipistrellus pipistrellus*). *Journal of Zoology*, 245: 222-228.
- OLSON, J. D. 1998. A digital model of pattern and productivity in an agroforestry landscape. *Landscape and Urban Planning*, 42: 169-189.
- ORESZCZYN, S. 2000. A systems approach to the research of people's relationships with English hedgerows. *Landscape and Urban Planning*, 50: 107-117.
- OSBORNE, J. L., MARTIN, A. P., SHORTALL, C. R., TODD, A. D., GOULSON, D., KNIGHT, M. E., HALE, R.J. & SANDERSON, R. A. 2008. Quantifying and comparing bumblebee nest densities in gardens and countryside habitats. *Journal of Applied Ecology*, 45: 784-792.
- OSBORNE, P. 1982. Some effects of Dutch Elm Disease on nesting farmland birds. *Bird Study*, 29: 2-16.
- OSBORNE, P. 1984. Bird numbers and habitat characteristics in farmland hedgerows. *Journal of Applied Ecology*, 21: 63-82.
- OUIN, A., & BUREL, F., 2002. Influence of herbaceous elements on butterfly diversity in hedgerow agricultural landscapes. *Agriculture, Ecosystems and Environment*, 93: 45-53.



- OUIN, A., AVIRON, S., DOVER, J. & BUREL, F. 2004. Complementation/supplementation of resources for butterflies in agricultural landscapes. *Agriculture, Ecosystems and Environment* 103: 473–479.
- OUIN, A., MARTIN, M., & BUREL, F. 2008. Agricultural landscape connectivity for the meadow brown butterfly (*Maniola jurtina*). *Agriculture, Ecosystems and Environment*, 124: 193-199.
- OWEN, J. A. 2003. Field studies on *Cryptocephalus coryli* (Linnaeus) (coleoptera: Chrysomelidae). *Entomologist's Gazette*, 54: 39-44.
- OWEN, J. A. 1999. Notes on the biology of *Cryptocephalus coryli* (Linnaeus) (Coleoptera: Chrysomelidae). *Entomologist's Gazette*, 50: 199-204.
- OWEN, J. A. 2000. Adult feeding and egg-laying in *Cryptocephalus coryli* (Linnaeus) (coleoptera: Chrysomelidae). *Entomologist's Gazette*, 51: 195-201.
- OWEN, J. A. 2002. Studies on the larval biology of *Cryptocephalus coryli* (Linnaeus) (coleoptera: Chrysomelidae). *Entomologist's Gazette*, 53: 57-68.
- OWEN, J. A. 2003. Studies on the life-history of *Cryptocephalus nitidulus* Fabricius, 1787 (coleoptera: Chrysomelidae). *Entomologist's Gazette*, 54: 255-266.
- OZANNE, C.M.P., FOGGO, A., HAMBLER, C. & SPEIGHT, M.R. 1997. The significance of edge-effects in the management of forests for invertebrate biodiversity. In Stork, N., Adis, J. & Didham, R., (eds), *Canopy Arthropods*. pp. 534–550. Chapman & Hall, London.
- PALETTA, A., CHINCARINI, M., & TOSI, V. 2008. An analysis of the variability of deadwood volumes found in hedgerows in the venet region (ITALY). [Une analyse de la variabilitÉ des volumes de bois mort observée dans les haies de vénétie (Italie)]. *Revue Forestiere Francaise*, 60: 437-450.
- PARISH, T., LAKHANI, K. H., & SPARKS, T. H. 1994. Modelling the relationship between bird population variables and hedgerow and other field margin attributes. I. Species richness of winter, summer and breeding birds. *Journal of Applied Ecology*, 31: 764-775.
- PARISH, T., LAKHANI, K. H., & SPARKS, T. H. 1995. Modelling the relationship between bird population variables and hedgerow, and other field margin attributes. II. Abundance of individual species and of groups of similar species. *Journal of Applied Ecology*, 32: 362-371.
- PEACH, W. J., DENNY, M., COTTON, P. A., HILL, I. F., GRUAR, D., BARRITT, D., IMPEY, A. & MALLORD, J. 2004. Habitat selection by song thrushes in stable and declining farmland populations. *Journal of Applied Ecology*, 41: 275-293.
- PENG, R. K., SUTTON, S. L., & FLETCHER, C. R. 1992. Spatial and temporal distribution patterns of flying Diptera. *Journal of Zoology*, 228: 329-340.
- PEREIRA, M., & RODRÍGUEZ, A. 2010. Conservation value of linear woody remnants for two forest carnivores in a Mediterranean agricultural landscape. *Journal of Applied Ecology*, 47: 611-620.
- PERKINS, A. J., WHITTINGHAM, M. J., MORRIS, A. J., & BRADBURY, R. B. 2002. Use of field margins by foraging yellowhammers *Emberiza citrinella*. *Agriculture, Ecosystems and Environment*, 93:413-420.
- PETIT, S., & BUREL, F. 1998. Connectivity in fragmented populations: *Abax parallelepipedus* in a hedgerow network landscape. [CONNECTIVITE BIOLOGIQUE ET POPULATIONS FRAGMENTEES: ABAX PARALLELEPIPEDUS DANS LES RESEAUX BOCAGERS]. *Comptes Rendus De l'Academie Des Sciences - Serie III*, 321: 55-61.
- PETIT, S., & BUREL, F. 1998. Effects of landscape dynamics on the metapopulation of a ground beetle (Coleoptera, carabidae) in a hedgerow network. *Agriculture, Ecosystems and Environment*, 69: 243-252.
- PETIT, S., & USHER, M. B. 1998. Biodiversity in agricultural landscapes: The ground beetle communities of woody uncultivated habitats. *Biodiversity and Conservation*, 7: 1549-1561.
- PIPER, R. 2002. The rare species of UK *Cryptocephalus*: the current state of knowledge and a possible way forward. Chapter 3 – PhD thesis 53pp



- POCOCK, M.O., EVANS, D.M. & MEMMOTT, J. 2012. The Robustness and Restoration of a Network of Ecological Networks. *Science*, 335: 973-977.
- POLLARD, E. 1968. Biological effects of shelter - interrelations between hedge and crop invertebrate faunas. In: Hooper, M.D. & Holdgate, M.W. (Eds.), *Hedges and Hedgerow Trees*. pp. 39-46. Monks Wood Experimental Station Symposium No.4, The Nature Conservancy, Peterborough.
- POLLARD, K. A., & HOLLAND, J. M. 2006. Arthropods within the woody element of hedgerows and their distribution pattern. *Agricultural and Forest Entomology*, 8: 203-211.
- PYWELL, R. F., JAMES, K. L., HERBERT, I., MEEK, W. R., CARVELL, C., BELL, D., & SPARKS, T. H. 2005. Determinants of overwintering habitat quality for beetles and spiders on arable farmland. *Biological Conservation*, 123: 79-90.
- PYWELL, R.F., MEEK, W.M., CARVELL, C., HULMES, L. & NOWAKOWSKI, M. 2007. The Buzz project: biodiversity enhancement on arable land under the new agri-environment schemes. *Aspects of Applied Biology*, 81: 61-68.
- PYWELL, R.F., WARMAN, E.A., CARVELL, C., SPARKS, T.H., DICKS, L.V., BENNETT, D., WRIGHT, A., CRITCHLEY, C.N.R. & SHERWOOD, A. 2005. Providing foraging resources for bumblebees in intensively farmed landscapes. *Biological Conservation*, 121: 479-494.
- PYWELL, R.F., WARMAN, E.A., HULMESA, L., HULMES, S., NUTTALLA, P., SPARKS, T.H., CRITCHLEY, C.N.R. & SHERWOOD, A. 2006. Effectiveness of new agri-environment schemes in providing foraging resources for bumblebees in intensively farmed landscapes. *Biological Conservation*, 129: 192-206.
- RANDS, M.R.W. 1986. Effect of hedgerow characteristics on partridge breeding densities. *Journal of Applied Ecology*, 23: 479-487.
- RAOUL, F., DEFAUT, R., MICHELAT, D., MONTADERT, M., PÉPIN, D., QUÉRÉ, J. P., TISSOT, B., DELATTRE P. & GIRAUDOUX, P. 2001. Landscape effects on the population dynamics of small mammal communities: A preliminary analysis of prey-resource variations. *Revue d'Ecologie (La Terre Et La Vie)*, 56: 339-352.
- READING, C.J & JOFRÉ, G.M. 2009. Habitat selection and range size of grass snakes *Natrix natrix* in an agricultural landscape in southern England. *Amphibia-Reptilia*, 30: 379-388.
- READING, C.J. & DAVIES C.L. 1996. Predation by grass snakes (*Natrix natrix*) at a site in southern England. *Journal of Zoology*, 239: 73-82.
- RICH, T. C. G., CLEMENTS, D. K., LEWIS, J., & MOORE, L. 2000. A comparison of four methods used to survey hedgerows: The Cardiff hedgerow survey 1998. *Journal of Environmental Management*, 60: 91-100.
- RICHARDSON, P. & FWAG, Undated. Bats and farms. FWAG Technical leaflet 2pp.
- RINK, M., & SINSCH, U. 2007. Radio-telemetric monitoring of dispersing stag beetles: Implications for conservation. *Journal of Zoology*, 272: 235-243.
- RINK, M., & SINSCH, U. 2011. Warm summers negatively affect duration of activity period and condition of adult stag beetles (*Lucanus cervus*). *Insect Conservation and Diversity*, 4: 15-22.
- RIORDAN, P., LLOYD, A. & MACDONALD, D.W. 2009. Do harvest mouse nest survey results predict population size? *Report to People's Trust for Endangered Species*. 10pp.
- ROBINSON, M.F. & STEBBINGS, R.E. 1997. Home range and habitat use by the serotine bat, *Eptesicus serotinus*, in England. *Journal of Zoology* 243: 117-136.
- ROBINSON, R. A., & SUTHERLAND, W. J. 1999. The winter distribution of seed-eating birds: Habitat structure, seed density and seasonal depletion. *Ecography*, 22: 447-454.
- ROBINSON, R. A., & SUTHERLAND, W. J. 2002. Post-war changes in arable farming and biodiversity in Great Britain. *Journal of Applied Ecology*, 39: 157-176.



- ROTHERAY, G. E. 1991. Larval stages of 17 rare and poorly known British hoverflies (Diptera: Syrphidae). *Journal of Natural History*, 25: 945-969.
- RUSS, J. M., & MONTGOMERY, W. I. 2002. Habitat associations of bats in Northern Ireland: Implications for conservation. *Biological Conservation*, 108:-58.
- RUSSO, D., JONES, G., & MIGLIOZZI, A. 2002. Habitat selection by the Mediterranean horseshoe bat, *Rhinolophus euryale* (chiroptera: Rhinolophidae) in a rural area of southern Italy and implications for conservation. *Biological Conservation*, 107: 71-81.
- SARLÖV HERLIN, I. L., & FRY, G. L. A. 2000. Dispersal of woody plants in forest edges and hedgerows in a southern Swedish agricultural area: The role of site and landscape structure. *Landscape Ecology*, 15: 229-242.
- SARLOV-HERLIN, I.L. 2004. Landscape ecology of woody tree and scrub species in hedgerows and woodland edges. Assumptions on relations between landscape composition and species occurrence, based on a field study. In: Smithers, R. (Ed.). *Landscape Ecology of Trees and Forests*: pp. 332-335 Proceedings of the 12th Annual IALE (UK) Conference, Cirencester, June 2004. pp. 332-335
- SCHIPPERS, P., GRASHOF-BOKDAM, C. J., VERBOOM, J., BAVECO, J. M., JOCHEM, R., MEEUWSEN, H. A. M., & VAN ADRICHEM, M. H. C. 2009Sacrificing patches for linear habitat elements enhances metapopulation performance of woodland birds in fragmented landscapes. *Landscape Ecology*, 24: 1123-1133.
- SCHOFIELD, H., MESSENGER, J., BIRKS, J. & JERMYN, D. 2002. *Foraging and roosting behaviour of lesser horseshoe bats at the Ciliau, Radnor*. The Vincent Wildlife Trust, Herefordshire, UK.25pp.<http://www.swild.ch/rhinolophus/Ciliau%20Report.pdf> (Accessed 18 February 2013).
- SHREEVE, T.G. & DENNIS, R.L.H. 2011. Landscape scale conservation: resources, behaviour, the matrix and opportunities. *Journal of Insect Conservation*, 15:179-188. doi:10.1007/s10841-010-9336-9
- SCHULZ, B., EHLERS, S., LANG, J. & BÜCHNER, S. 2012. Hazel dormice in roadside habitats. *Peckiana*, 8: 49-55.
- SHERIDAN, H., FINN, J. A., & O'DONOVAN, G. 2009. Botanical rejuvenation of field margins and benefits for invertebrate fauna on a drystock farm in County Longford. *Biology and Environment*, 109: 95-106.
- SIRIWARDENA, G.M., COOKE, I.R. & SUTHERLAND, W.J. 2012. Landscape, cropping and field boundary influences on bird abundance. *Ecography*, 35: 162-173.
- SMITH, J., POTTS, S., & EGGLETON, P. 2008. The value of sown grass margins for enhancing soil macrofaunal biodiversity in arable systems. *Agriculture, Ecosystems and Environment*, 127: 119-125.
- SMITH, J., POTTS, S.G., WOODCOCK, B.A. & EGGLETON, P. 2008. Can arable field margins be managed to enhance their biodiversity, conservation and functional value for soil macrofauna? *Journal of Applied Ecology*, 45: 269–278.
- SMITH, P.G. & RACEY, P.A. 2005. The itinerant Natterer: physical and thermal characteristics of summer roosts of *Myotis nattereri* (Mammalia: Chiroptera). *Journal of Zoology* 266, 171-180.
- SPARKS, T. H. & CROXTON, P.J. 2007. The influence of timing and frequency of hedgerow cutting on hawthorn flowering and berry yields: preliminary results. *Aspects of Applied Biology*, 82: 103 – 106.
- SPARKS, T. H., & PARISH, T. 1995. Factors affecting the abundance of butterflies in field boundaries in Swavesey Fens, Cambridgeshire, UK. *Biological Conservation*, 73: 221-227.
- SPARKS, T. H., PARISH, T., & HINSLY, S. A. 1996. Breeding birds in field boundaries in an agricultural landscape. *Agriculture, Ecosystems and Environment*, 60: 1-8.
- SPEIGHT, M.C.D. 2001. Farms as biogeographical units:2. The potential role of different parts of the case-study farm in maintaining its present fauna of Sciomyzidae and Syrphidae (Diptera). *Bulletin of the Irish Biogeographical Society*, 25: 248-278.
- SPELLERBERG, I. F. & GAYWOOD, M.J. 1993. *Linear features: linear habitats & wildlife corridors*. English Nature Research Report 60. 74pp.



- STALEY, J. T., SPARKS, T. H., CROXTON, P. J., BALDOCK, K. C. R., HEARD, M. S., HULMES, S., PEYTON' J., AMY, S.R. & PYWELL, R. F. 2012. Long-term effects of hedgerow management policies on resource provision for wildlife. *Biological Conservation*, 145: 24-29.
- STOKES, J. & HAND, K. 2004. *The Hedge Tree Handbook*. The Tree Council, London.
- TATTERSALL, F. H., MACDONALD, D. W., HART, B. J., JOHNSON, P., MANLEY, W., & FEBER, R. 2002. Is habitat linearity important for small mammal communities on farmland? *Journal of Applied Ecology*, 39: 643-652.
- THOMAES, A., KERVYN, T., & MAES, D. 2008. Applying species distribution modelling for the conservation of the threatened saproxylic stag beetle (*Lucanus cervus*). *Biological Conservation*, 141: 1400-1410.
- THOMAES, A., KERVYN, T., BECK, O., & CAMMAERTS, R. 2008. Distribution of *Lucanus cervus* (coleoptera: Lucanidae) in Belgium: Surviving in a changing landscape. *Revue d'Ecologie (La Terre Et La Vie)*, 63: 147-152.
- THOMAS, C. F. G., & MARSHALL, E. J. P. 1999. Arthropod abundance and diversity in differently vegetated margins of arable fields. *Agriculture, Ecosystems and Environment*, 72: 131-144.
- THOMAS, C. F. G., GREEN, F., & MARSHALL, E. J. P. 1998. Distribution, dispersal and population size of the ground beetles, *Pterostichus melanarius* (Illiger) and *Harpalus rufipes* (Degeer) (Coleoptera, Carabidae), in field margin habitats. *Biological Agriculture and Horticulture*, 15: 337-352.
- THOMAS, C. F. G., PARKINSON, L., & MARSHALL, E. J. P. 1998. Isolating the components of activity-density for the carabid beetle *Pterostichus melanarius* in farmland. *Oecologia*, 116: 103-112.
- THOMAS, C. F. G., PARKINSON, L., GRIFFITHS, G. J. K., GARCIA, A. F., & MARSHALL, E. J. P. 2001. Aggregation and temporal stability of carabid beetle distributions in field and hedgerow habitats. *Journal of Applied Ecology*, 38: 100-116.
- THOMAS, C.D. 2005. Monitoring change in the abundance and distribution of insects using butterflies and other indicator groups. *Philosophical Transactions of the Royal Society of London Series B*, 360: 339-357.
- THOMAS, J. A., BOURN, N.A.D., CLARKE, R.T., STEWART, K.E., SIMCOX, D.J., PEARMAN, G.S., CURTIS, R. & GOODGER, B. 2001. The quality and isolation of habitat patches both determine where butterflies persist in fragmented landscapes. *Proc. R. Soc. Lond. B 7 September 2001 vol. 268 no. 1478 1791-1796*
- THOMAS, S. R., GOULSON, D., & HOLLAND, J. M. 2001. Resource provision for farmland gamebirds: The value of beetle banks. *Annals of Applied Biology*, 139: 111-118.
- THOMAS, S. R., NOORDHUIS, R., HOLLAND, J. M. & GOULSON, D. 2002. Botanical diversity of beetle banks. Effects of age and comparison with conventional arable field margins in southern UK. *Agriculture, Ecosystems and Environment*, 93: 403-412.
- TISCHENDORF, L., IRMLER, U., & HINGST, R. 1998. A simulation experiment on the potential of hedgerows as movement corridors for forest carabids. *Ecological Modelling*, 106: 107-118.
- USIETA, H.O., MANU' S.A. & OTTOSSON' U. 2013. Farmland conservation in West Africa: how do hedgerow characteristics affect bird species richness? *Bird Study*, 60: DOI10.1080/00063657.2012.758226
- VAN LANGEVELDE, F., & GRASHOF-BOKDAM, C. J. 2011. Modelling the effect of intersections in linear habitat on spatial distribution and local population density. *International Journal of Geographical Information Science*, 25: 367-378.
- VARCHOLA, J. M., & DUNN, J. P. 2001. Influence of hedgerow and grassy field borders on ground beetle (Coleoptera: Carabidae) activity in fields of corn. *Agriculture, Ecosystems and Environment*, 83: 153-163.
- VARET, M., PÉTILLON, J., & BUREL, F. 2011. Comparative responses of spider and carabid beetle assemblages along an urban-rural boundary gradient. *Journal of Arachnology*, 39: 236-243.
- VERA, F.W.M. 2000. *Grazing Ecology and Forest History*. CABI, Wallingford, UK. 32pp.
- VERBOOM, B. & HUITEMA, H. 1997. The importance of linear landscape elements for the pipistrelle *Pipistrellus pipistrellus* and the serotine bat *Eptesicus serotinus*. *Landscape Ecology*, 12: 117-125.



- VERHEYEN, K., GUNTENSPERGEN, G. R., BEISBROUCK, B., & HERMY, M. 2003. An integrated analysis of the effects of past land use on forest herb colonization at the landscape scale. *Journal of Ecology*, 91: 731-742.
- VICKERY, J. A., FEEBER, R. E., & FULLER, R. J. 2009. Arable field margins managed for biodiversity conservation: A review of food resource provision for farmland birds. *Agriculture, Ecosystems and Environment*, 133: 1-13.
- VIGNON, V. & ORABI, P. 2003. Exploring the hedgerows network in the west of France for the conservation of saproxylic beetles (*Osmoderma eremita*, *Gnorimus variabilis*, *Lucanus cervus*, *Cerambyx cerdo*). *Proceedings of the second pan-European conference on Saproxylic Beetles* pp 1-3. People's Trust for Endangered Species.
- VINCENT, S., NEMOZ, M., & AULAGNIER, S. 2011. Activity and foraging habitats of *Miniopterus schreibersii* (chiroptera, miniopteridae) in southern France: Implications for its conservation. [Attività e habitat di caccia di *Miniopterus schreibersii* (Chiroptera, Miniopteridae) nel sud della Francia: Implicazioni per la sua conservazione]. *Hystrix*, 22: 57-72.
- WALKER, M. P., DOVER, J. W., HINSLEY, S. A., & SPARKS, T. H. 2005. Birds and green lanes: Breeding season bird abundance, territories and species richness. *Biological Conservation*, 126: 540-547.
- WALKER, M.P., DOVER, J.W., SPARKS, T.H. & HINSLEY, S.A. 2006. Hedges and green lanes: Vegetation composition and structure. *Human Exploitation and Biological Conservation*, 3:255-270.
- WALSH A.L., & HARRIS, S. 1996. Foraging habitat preferences of vespertilionid bats in Britain. *Journal of Applied Ecology*, 33: 508-518.
- WALTHER-HELLWIG, K., & FRANKL, R. 2000. Foraging habitats and foraging distances of bumblebees, *Bombus* spp. (Hym., apidae), in an agricultural landscape. *Journal of Applied Entomology*, 124: 299-306.
- WARING, P. 2004. Successes in conserving the Barberry Carpet moth *Pareulype berberata* (D. & S.) (Geometridae) in England. *Journal of Insect Conservation* 8: 167-171.
- WATT, T.A. & BUCKLEY G.P. 1994. *Hedgerow Management & Nature Conservation*. Wye College, University of London, 4-5 Sept 1992. Wye College, Ashford.
- WEBB, J.R., DREWITT, A.L., & MEASURES, G.H. 2009. *Managing for species: Integrating the needs of England's priority species into habitat management. Part 2 Annexes*. Natural England Research Reports, Number 024. 94pp.
- WEBB, J.R., DREWITT, A.L., & MEASURES, G.H. 2010. *Managing for species: Integrating the needs of England's priority species into habitat management. Part 1 Report*. Natural England Research Reports, Number 024. 146pp.
- WEHLING, S. & DIEKMANN, M. 2009. Importance of hedgerows as habitat corridors for forest plants in agricultural landscapes. *Biological Conservation*, 142: 2522-2530.
- WERMELINGER, B., FLÜCKIGER, P.F., OBRIST, M.K. & DUELLI, P. 2007. Horizontal and vertical distribution of saproxylic beetles (Col., Buprestidae, Cerambycidae, Scolytinae) across sections of forest edges. *Journal of Applied Entomology*, 131: 104-114.
- WHITTINGHAM, M. J., BRADBURY, R. B., WILSON, J. D., MORRIS, A. J., PERKINS, A. J., & SIRIWARDENA, G. M. 2001. Chaffinch *Fringilla coelebs* foraging patterns, nestling survival and territory distribution on lowland farmland. *Bird Study*, 48: 257-270.
- WHITTINGHAM, M. J., SWETNAM, R. D., WILSON, J. D., CHAMBERLAIN, D. E., & FRECKLETON, R. P. 2005. Habitat selection by yellowhammers *Emberiza citrinella* on lowland farmland at two spatial scales: Implications for conservation management. *Journal of Applied Ecology*, 42: 270-280.
- WHITTINGHAM, M.J. & EVANS, K.L. 2004. The effects of habitat structure on predation risk of birds in agricultural landscapes. *Ibis*, 146: 210-220.
- WICKRAMASINGHE, L. P., HARRIS, S., JONES, G., & VAUGHAN, N. 2003. Bat activity and species richness on organic and conventional farms: Impact of agricultural intensification. *Journal of Applied Ecology*, 40: 984-993.
- WILLIAMS, M. 2008. Hedgerows For Hairstreaks? A Farm Hedgerow Survey Report. *Butterfly Conservation*, 13pp.



- WILSON, J. D., MORRIS, A. J., ARROYO, B. E., CLARK, S. C., & BRADBURY, R. B. 1999. A review of the abundance and diversity of invertebrate and plant foods of granivorous birds in northern Europe in relation to agricultural change. *Agriculture, Ecosystems and Environment*, 75: 13-30.
- WINSPEAR, R. & DAVIES, G. 2005. *A management guide to birds of lowland farmland*. RSPB, Sandy, 169pp.
- WISLER, C., ULRICH HOFER, U. & ARLETTAZ, R.L. 2008. Snakes and Monocultures: Habitat Selection and Movements of Female Grass Snakes (*Natrix natrix* L.) in an Agricultural Landscape. *Journal of Herpetology*, 42: 337–346.
- WOODCOCK, B.A., WESTBURY, D.B., POTTS, S.G., HARRIS, S.J. & BROWN, V.K. 2005. Establishing field margins to promote beetle conservation in arable farms. *Agriculture, Ecosystems and Environment*, 107: 255–266.
- WOODCOCK, B.A., WESTBURY, D.B., TSCHÉULIN, T., HARRISON-CRIPPS, J., HARRIS, S.J., RAMSEY, A.J., BROWN, V.K., POTTS, S.G., 2008. Effects of seed mixture and management on beetle assemblages of arable field margins. *Agriculture, Ecosystems & Environment*, 125: 246–254.
- WOODCOCK B.A., VOGIATZAKIS I.N., WESTBURY D.B., LAWSON C.S., EDWARDS A.R., BROOK A.J., HARRIS S.J., LOCK K.A., MACZEY N. & MASTERS G. 2010. The role of management and landscape context in the restoration of grassland phytophagous beetles. *Journal of Applied Ecology*, 47, 366-376.
- WOODCOCK, B.A., BULLOCK, J.M., MORTIMER, S.R., BRERETON, T., REDHEAD, J.W., THOMAS, J.A. & PYWELL, R.F. 2012. Identifying time lags in the restoration of grassland butterfly communities: A multi-site assessment. *Biological Conservation*, 155: 50–58.
- WOLTON, R. J. 2008. Where dormice nest in hedges. The Dormouse Monitor, Spring 2008 pp6-7.
- WOLTON, R. J. 2009. Hazel dormouse *Muscardinus avellanarius* (L.) nest site selection in hedgerows. *Mammalia*, 73: 7-12.
- WOLTON, R. J. 2009. *UK Biodiversity Action Plan: Priority species linked to hedgerows*. Report by Hedgelink, for Defra
- WOLTON, R.J. & VERGETTE, M. 2012. The diversity of life in a single hedge. In Dover, J.W., (ed.) *Hedgerow Futures* pp. 98-104.
- WRATTEN, S.D., BOWIE, M. H., HICKMAN, J. M., EVANS, A. M., SEDCOLE, J. R., & TYLIANAKIS, J. M. 2003. Field boundaries as barriers to movement of hover flies (Diptera: Syrphidae) in cultivated land. *Oecologia*, 134: 605-611.
- WRATTEN, S.D., GILLESPIE, M., DECOURTYE, A., MADER, E., & DESNEUX, N. 2012. Pollinator habitat enhancement: Benefits to other ecosystem services. *Agriculture, Ecosystems and Environment*, 159: 112-122.
- ZEALE, M. R.K., DAVIDSON-WATTS, I., & JONES, G. 2012. Home range use and habitat selection by Barbastelle bats (*Barbastella barbastellus*): Implications for conservation. *Journal of Mammalogy*, 93: 1110-1118.

